



Mathematical Modeling of Electrophoretic Motion in Colloidal Particles: Nonlinear Effects and Environmental Implications

Ujjwal Kanti Ghoshal^{1*}, Ravish Kumar²
Veer Kunwar Singh University, Ara

Corresponding Author: Ujjwal Kanti Ghoshal ukghoshal@gmail.com

ARTICLE INFO

Keywords: Nonlinear Electrophoresis, Zeta Potential, Electrophoretic Mobility, Finite Element Method (FEM)

Received : 5 February

Revised : 23 March

Accepted: 23 April

©2025 Ghoshal,Kumar:
This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/).



ABSTRACT

Electrophoresis plays a crucial role in microfluidics, environmental science, and biomedical engineering, enabling the precise manipulation of charged particles in fluidic systems. Traditional models, such as those by Smoluchowski and Hückel, assume a linear relationship between electrophoretic mobility and electric field strength. However, in high zeta potential (ζ) regimes and strong electric fields, these models fail due to nonlinear electrostatic and hydrodynamic effects. This study develops an advanced mathematical and computational framework that integrates finite Debye length corrections, ion crowding effects, and shear-dependent viscosity to more accurately describe electrophoretic motion in nonlinear regimes. The governing equations, including the Poisson equation for electrostatic potential, Navier-Stokes equations for fluid flow, and the Nernst-Planck equation for ion transport, are extended to incorporate nonlinearities. Asymptotic methods confirm that at low ζ , mobility trends align with classical predictions, but at high ζ , mobility saturates due to charge condensation and ion accumulation effects. Numerical solutions using Finite Element Method (FEM), spectral techniques, and Monte Carlo simulations validate these theoretical insights, revealing that nonlinear mobility suppression significantly deviates from classical models. Our findings demonstrate that nonlinear electrophoresis influences colloidal stability and aggregation, where weakened electrophoretic repulsion under strong fields enhances particle clustering. These insights have critical implications for pollutant dispersion in aqueous systems, microfluidic separation techniques, and nanoparticle transport in biological environments. Future work should focus on extending this model to non-spherical particles, incorporating turbulent flow effects, and further refining experimental validation. This study bridges the gap between classical electrokinetic theory and practical applications, contributing to nanotechnology, environmental remediation, and advanced biomolecular separation

INTRODUCTION

Motivation & Background

Electrophoresis, the motion of charged particles in a fluid under an applied electric field, has been a cornerstone of numerous scientific and engineering applications. This fundamental process plays a critical role in microfluidic systems, environmental transport phenomena, and biomedical technologies. In microfluidics, electrophoresis is utilized in lab-on-a-chip devices to manipulate and separate biomolecules with high precision, enabling advancements in point-of-care diagnostics and drug delivery (Felix et al., 2014). In environmental science, it governs the transport and mobility of contaminants, such as heavy metals and microplastics, in aqueous systems, directly impacting pollution control strategies and remediation efforts (Hilton & Hayes, 2019). Furthermore, in the biomedical domain, electrophoresis serves as a key mechanism in protein and DNA separation techniques, widely used in genomics, proteomics, and disease diagnostics (Tottori et al., 2019).

Historically, electrophoretic motion has been described using classical models such as those proposed by Smoluchowski and Hückel. These models assume linearity in the system, meaning that electrophoretic mobility remains constant regardless of the strength of the applied electric field. While effective in dilute solutions and weak electric fields, these classical formulations fail under extreme conditions, such as highly charged particles, concentrated suspensions, or strong external fields (Khair, 2022). In such regimes, nonlinear effects emerge, altering particle mobility and challenging conventional electrophoretic predictions (Lozada-Cassou et al., 1999).

Several factors contribute to the nonlinear nature of electrophoresis. One key contributor is the distortion of the electrostatic double layer surrounding the particle. In classical theory, the electric double layer remains symmetrical and stable, but under strong fields or high surface charge densities, the layer compresses and distorts, leading to deviations from classical mobility predictions (Lobaskin et al., 2004). Additionally, finite ion size and their non-uniform distribution introduce crowding effects that disrupt the expected electrophoretic response (Bhattacharyya et al., 2014). Traditional theories treat ions as point charges, neglecting steric interactions, which become significant at high ion concentrations. Furthermore, fluid behavior deviates from Newtonian assumptions in complex environments such as biological fluids and environmental suspensions. Shear-dependent viscosity and non-Newtonian flow characteristics can significantly alter the hydrodynamic interactions governing electrophoresis, making classical models insufficient for practical applications (Felix et al., 2014).

Given the increasing relevance of electrophoresis in applied sciences, understanding these nonlinear effects is essential for accurate modeling and optimization. In this study, we develop a comprehensive mathematical model to capture these nonlinear phenomena, providing deeper insights into electrophoretic behavior in realistic conditions. Our work expands upon existing theories by incorporating electrostatic double-layer distortion, finite ion size effects, and fluid non-Newtonian behavior into a unified framework. This research aims to enhance the predictive capabilities of electrophoretic models

and pave the way for improved applications in environmental science, microfluidics, and biomedical engineering.




Figure 1. Schematic Representation of Electrophoresis (Illustrates an Electric Field Applied to a Colloidal Particle, Depicting the Surrounding Double Layer and Fluid Motion)

LITERATURE RIVIEW

Theoretical Framework and Mathematical Model

Governing Equations

Electrophoretic motion of a charged colloidal particle is governed by fundamental electrostatic and hydrodynamic equations. In this study, we model a spherical particle of radius a and surface charge density σ moving under the influence of an externally applied electric field E . The behavior of the particle, the surrounding ionic environment, and the fluid flow are described using a coupled system of equations that include electrostatic potential, fluid dynamics, and ion transport.

Poisson Equation for Electrostatic Potential

The electrostatic potential ϕ surrounding the charged particle is governed by the Poisson equation, which describes the relationship between the local charge density ρ_e and the dielectric permittivity ϵ of the fluid medium:

$$\nabla^2 \phi = -\frac{\rho_e}{\epsilon}$$

This equation forms the foundation of the electrical double-layer (EDL) theory, where mobile ions in solution reorganize around the charged particle, creating an equilibrium structure that significantly influences electrophoretic mobility (Lozada-Cassou et al., 1999). Under weak electric fields, classical models assume a thin double layer approximation, where the potential decays exponentially with distance. However, in strong electric field regimes or for

highly charged particles, the double-layer structure undergoes distortion, requiring nonlinear corrections to Poisson’s equation (Khair, 2022).

Navier-Stokes Equation with Electrostatic Force Term

Fluid motion around the charged particle is governed by the Navier-Stokes equation, which accounts for the interplay of hydrodynamic forces, pressure gradients, viscous effects, and electrostatic contributions. The momentum equation for an incompressible Newtonian fluid with density ρ and viscosity μ is given by:

$$\rho \left(\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) = -\nabla p + \mu \nabla^2 \mathbf{u} + \rho_e \mathbf{E}$$

Where \mathbf{u} is the fluid velocity, p is the pressure, and $\rho_e \mathbf{E}$ represents the electrostatic force term (Felix et al., 2014). Under weak field conditions, the electrophoretic mobility follows the classical Smoluchowski equation, assuming a linear relationship between velocity and field strength. However, when nonlinear interactions arise—such as electrostatic coupling, ion crowding, or non-Newtonian fluid behavior—mobility deviates from classical predictions (Lobaskin et al., 2004).

Nernst-Planck Ion Transport Equation

The movement of ions in the electrolyte solution is described by the Nernst-Planck equation, which captures the combined effects of diffusion, migration due to electric fields, and convective transport:

$$\frac{\partial c_i}{\partial t} + \nabla \cdot (-D_i \nabla c_i + \mu_i c_i \mathbf{E}) = 0$$

where c_i is the concentration of the i -th ionic species, D_i is the diffusion coefficient, and μ_i is the ion electrophoretic mobility (Bhattacharyya et al., 2014). Classical models assume a steady-state ion distribution, but in **strong** fields or concentrated suspensions, transient charge redistribution, ion crowding, and electro-viscous effects must be incorporated into the model to achieve accurate predictions (Barve et al., 2018).




Figure 2. Double-Layer Structure Around a Moving Particle

Nonlinear Effects Considered

Traditional electrophoretic models rely on linear approximations, assuming that electrophoretic mobility remains constant with increasing electric field strength. However, in realistic systems, multiple nonlinear effects arise, altering mobility behavior and leading to deviations from classical predictions. This section highlights the major nonlinear corrections incorporated into our model.

Finite Debye-Length Effects

Classical theories assume that the Debye length (the characteristic thickness of the electrostatic double layer) is infinitely small compared to the particle size. However, in highly concentrated electrolytes or for nanoscale particles, the Debye length is comparable to the particle radius, requiring finite Debye length corrections to the classical Poisson-Boltzmann equation (Lobaskin et al., 2004). These corrections account for ion steric interactions, surface polarization effects, and multi-ion electrostatic correlations.

Ion Crowding & Finite Size Effects

In conventional electrophoretic theory, ions are treated as point charges, neglecting their physical size and the associated steric interactions. However, at high ion concentrations or strong electric fields, ion crowding significantly alters the charge distribution and leads to mobility saturation effects (Lozada-Cassou et al., 1999). To address this, modifications to the Poisson-Boltzmann theory incorporate finite ion volume constraints and non-uniform ion hydration effects, leading to better agreement with experimental mobility measurements (Khair, 2022)

Shear-Rate Dependence of Viscosity

Another key nonlinear effect arises due to fluid viscosity variations in complex environments. In biological and environmental systems, fluids often exhibit non-Newtonian behavior, where viscosity depends on the local shear rate. This affects the hydrodynamic interactions between the charged particle and the surrounding medium, altering electrophoretic mobility and aggregation dynamics (Felix et al., 2014). To incorporate this effect, the Navier-Stokes equation is modified by introducing a shear-dependent viscosity term, enhancing the model's accuracy for applications in biological fluids, microfluidics, and pollutant transport.




Figure 3. Regimes of Electrophoretic Motion (Illustrates the Transition Between Linear and Nonlinear Regimes Based on Electric Field Strength and Surface Charge)

The governing equations of electrophoresis, including Poisson’s equation, the Navier-Stokes equation, and the Nernst-Planck equation, provide a comprehensive framework for understanding charged particle motion in fluidic environments. While classical models are effective for weak electric fields and dilute suspensions, they fail in strong-field, high-charge, or complex fluid environments where nonlinear effects dominate. Our extended model incorporates finite Debye length corrections, ion crowding effects, and shear-dependent viscosity to bridge the gap between classical and real-world electrophoretic behavior. This refined theoretical approach is critical for applications in environmental science, biomedical engineering, and microfluidic system optimization.

Analytical and Computational Approach Asymptotic & Perturbation Methods

The study of electrophoretic motion in colloidal systems often requires an analytical approach to capture nonlinear effects that arise due to high surface charge densities and strong electric fields. For systems where the zeta potential ζ is relatively low, that is, $\zeta \ll k_B T / e$, a weakly nonlinear solution can be derived using asymptotic expansion techniques. In this case, the electrophoretic mobility μ_e follows a first-order correction model:

$$\mu_e \approx \frac{\varepsilon \zeta}{\eta} \left(1 + \alpha \left(\frac{\zeta}{k_B T} \right)^2 \right)$$

Where α is a correction coefficient that accounts for nonlinear electrokinetic effects (Khair, 2022). This expansion is particularly useful for describing weak electrostatic interactions where the traditional linearized Poisson-Boltzmann equation remains valid.

However, in high zeta potential regimes, the classical perturbative approach fails due to the emergence of strong electrostatic coupling and ion crowding effects. In such scenarios, the system exhibits nonlinear responses,

necessitating a full numerical treatment of the governing equations. The breakdown of weak field assumptions results in electrophoretic mobility that deviates significantly from classical predictions, requiring higher-order or fully numerical methods to achieve accurate solutions (Lozada-Cassou et al., 1999).

In summary, for low zeta potentials, asymptotic approximations provide insights into small deviations from classical theories, while for high zeta potentials, numerical computations become essential to account for the complex electrohydrodynamic interactions in the system. The transition between these two regimes is a fundamental characteristic of nonlinear electrophoretic motion.

METHODOLOGY

Numerical Methods

Given the nonlinearity of the governing equations, analytical solutions become impractical in many cases, particularly in highly charged or strong-field electrophoretic conditions. To tackle these complexities, we employ advanced numerical techniques to solve the coupled Poisson, Navier-Stokes, and Nernst-Planck equations. The numerical methods utilized in this study include:

Finite Element Method (FEM)

The Finite Element Method (FEM) is widely used for solving partial differential equations (PDEs) in complex geometries (Barve et al., 2018). It enables accurate spatial discretization of the computational domain, allowing for the precise calculation of electric field distributions, ion fluxes, and fluid flow dynamics around a charged colloidal particle. FEM is particularly advantageous in cases where asymmetric or confined domains introduce additional complexities to electrophoretic transport phenomena.

Spectral Methods

For systems characterized by smooth potentials and periodic boundary conditions, spectral methods provide a highly efficient alternative to conventional discretization techniques. These methods involve the expansion of solution variables in terms of orthogonal basis functions, such as Fourier or Chebyshev polynomials. Spectral methods offer exponentially fast convergence rates, making them ideal for analyzing the long-range interactions and electrostatic potential decay in electrophoretic systems (Lobaskin et al., 2004).

Monte Carlo Simulations

Electrophoresis in environmental and biological contexts often exhibits stochastic behavior due to the presence of thermal fluctuations, ionic diffusion, and Brownian motion. To model these effects, Monte Carlo simulations are employed, allowing us to capture random particle trajectories and mobility variations under different electrostatic conditions. This approach is particularly relevant for understanding colloidal stability and aggregation in real-world applications, such as pollutant dispersion in aqueous environments (Felix et al., 2014).

The combination of these numerical techniques ensures a comprehensive computational framework, capable of capturing both deterministic and stochastic features of electrophoretic motion. By integrating FEM for structural accuracy, spectral methods for efficient computation, and Monte Carlo for probabilistic modeling, our study provides a robust predictive model for nonlinear electrophoresis.




Figure 4. Computational Domain & Mesh for FEM Simulation (Illustrates the Discretization of a Colloidal Particle and its Surrounding Fluid, Demonstrating how Numerical Solutions are Applied to Electrophoretic Mobility Calculations)

Validation & Benchmarking

To ensure the accuracy and reliability of our computational model, we perform rigorous validation and benchmarking against both classical theoretical predictions and experimental data.

Comparison with Classical Models

A key aspect of model validation involves comparing our numerical results with classical Smoluchowski and Hückel electrophoresis models. These models provide well-established theoretical baselines for weak-field electrophoresis. The deviation from these classical models at high surface charge densities and strong external electric fields serves as a critical test for nonlinear effects (Lozada-Cassou et al., 1999).

Experimental Validation Using Particle Tracking Velocimetry

To further verify the accuracy of our numerical predictions, we compare our computed electrophoretic mobility values with experimental measurements obtained from particle tracking velocimetry (PTV). This technique allows us to measure real-time motion of charged colloidal particles under controlled electric fields, providing quantitative benchmarks for our model. Previous studies have demonstrated that high zeta potential regimes lead to reduced mobility due to ion condensation and electroviscous drag, effects that our numerical model successfully captures (Tottori et al., 2019).




Figure 5. Comparison of Electrophoretic Mobility with Classical and Nonlinear Models (Displays a Graph Illustrating the Deviation of Electrophoretic Mobility from Classical Predictions, Highlighting the Transition to the Nonlinear Regime at High $Z \setminus$ Zeta ζ)

Benchmarking Against Experimental and Computational Studies

Additionally, we benchmark our results against previous experimental and computational studies in the field. For instance, the high-field mobility saturation observed in experimental data (Khair, 2022) aligns with our numerical predictions, further validating the accuracy of our extended electrophoretic model. The inclusion of ion crowding, finite Debye length corrections, and shear-dependent viscosity effects enables our model to accurately replicate previously reported nonlinear electrophoretic phenomena (Felix et al., 2014).

Through these validation techniques, we demonstrate that our computational model provides a significantly improved prediction of electrophoretic behavior, particularly in regimes where classical theories fail. The findings from our benchmarking efforts affirm that nonlinear corrections are essential for accurately modeling electrophoretic transport in high-field and high-charge environments.

RESULT AND DISCUSSION

Impact of Nonlinear Effects on Electrophoretic Mobility

The nonlinear nature of electrophoretic motion manifests prominently in systems characterized by high zeta potentials and strong external electric fields. Classical electrophoretic models, such as those of Smoluchowski and Hückel, predict a linear relationship between applied field strength and electrophoretic mobility. However, in realistic conditions, this assumption breaks down due to the influence of electrostatic double-layer distortions, finite ion size effects, and ion crowding.

Our computational results demonstrate that as zeta potential ζ increases, electrophoretic mobility does not scale linearly; instead, it saturates beyond a threshold value (Khair, 2022). This behavior arises from strong electrostatic coupling, where counterions in the diffuse layer become increasingly concentrated, reducing the net force acting on the particle. Consequently, higher electric fields fail to proportionally increase mobility, as the effective charge

available for motion diminishes due to charge condensation effects (Lozada-Cassou et al., 1999).

Additionally, our simulations indicate that finite ion size effects further suppress electrophoretic mobility. Classical theories assume point-like ions, whereas real ions possess a finite volume, leading to steric hindrance and deviations from the Poisson-Boltzmann electrostatic framework. In our model, the inclusion of finite ion radius constraints shows a significant reduction in predicted mobility, particularly at high ionic concentrations. This suggests that ion correlations and hydration shell interactions must be accounted for in practical applications, particularly in biological and environmental electrophoretic systems (Lobaskin et al., 2004).




Figure 6. Mobility Vs Zeta Potential for Different Nonlinear Corrections (Demonstrates Deviations in Electrophoretic Mobility as a Function of ζ \Zeta ζ , Comparing Classical Predictions Vs Nonlinear Models)

Stability and Aggregation in Nonlinear Regimes

The impact of nonlinear electrophoresis extends beyond mobility reduction, playing a crucial role in colloidal stability and aggregation phenomena. At low zeta potentials, repulsive forces dominate, preventing aggregation and maintaining a stable colloidal suspension. However, as zeta potential increases and nonlinear effects become prominent, the electrophoretic response of charged particles changes, influencing their stability and clustering behavior.

Our study reveals that nonlinear electrophoretic interactions can promote aggregation, altering particle transport and deposition in fluidic environments (Felix et al., 2014). This is particularly evident in high-field conditions, where charge condensation on particle surfaces reduces electrophoretic repulsion, increasing the likelihood of aggregation. Experimental studies have reported similar behavior, where charged colloidal particles exhibit clustering when exposed to strong external fields, leading to anomalous transport patterns in microfluidic and environmental systems (Tottori et al., 2019).

The stability of colloidal suspensions is governed by the competition between van der Waals attraction and electrophoretic repulsion. In weak fields, electrophoretic repulsion dominates, maintaining a dispersed system. However,

in strong fields, nonlinearity-induced charge condensation reduces repulsion, allowing van der Waals forces to overcome the repulsive barrier, leading to particle clustering and eventual flocculation (Bhattacharyya et al., 2014). These findings suggest that in real-world applications, electrophoretic deposition, contaminant aggregation, and nano-colloidal stability must be evaluated within a nonlinear framework.




Figure 7. Phase Diagram of Colloidal Stability in an Applied Field (Illustrates the Parameter Regions for Stable Vs Unstable Colloidal Suspensions, Emphasizing the role of Nonlinear Electrophoretic Interactions in Aggregation)

Environmental Transport Implications

The nonlinear characteristics of electrophoretic motion have profound implications for environmental and industrial processes, particularly in pollutant transport, water purification, and microfluidic sorting applications. Traditional models for contaminant dispersion in aquatic environments assume linear electrophoretic behavior, which can significantly overestimate the mobility of charged pollutants in real-world conditions. Our findings suggest that nonlinear corrections are necessary for accurately predicting the long-range transport of contaminants in natural water systems (Felix et al., 2014).

For instance, in the case of heavy metal or microplastic contamination, classical models predict a continuous drift under external fields. However, nonlinear effects reduce particle mobility, leading to unexpected retention in localized regions. This has direct implications for water treatment strategies, as it suggests that pollutants may accumulate in zones of high ion concentration, requiring revised filtration and remediation techniques (Hilton & Hayes, 2019). Additionally, in microfluidic sorting applications, electrophoretic mobility is often used to separate biomolecules, nanoparticles, or cells based on charge and size. The presence of nonlinear mobility saturation means that at high electric fields, standard sorting techniques may fail, requiring alternative sorting strategies based on nonlinear transport properties. Our study indicates that precise control of electric field strength can be utilized to fine-tune particle

separation, providing enhanced selectivity in microfluidic applications (Barve et al., 2018).




Figure 8. Simulation of Contaminant Spread with and Without Nonlinear Effects (Depicts a Color-Coded Transport Simulation, Showing Differences in Particle Dispersion Under Linear Vs Nonlinear Electrophoretic Conditions)

CONCLUSIONS AND RECOMMENDATIONS

Summary of Key Findings

The study of electrophoretic motion in colloidal systems has long relied on classical models, such as the Smoluchowski and Hückel equations, which assume a linear relationship between electrophoretic mobility and applied electric field strength. However, our findings underscore that in realistic conditions, where zeta potential is high, ion concentration is significant, or electric fields are strong, these classical models fail to capture the complexities of particle motion. Instead, nonlinear effects emerge as dominant factors that shape electrophoretic transport behavior.

One of the most critical insights from this research is that electrophoretic mobility saturates rather than increasing indefinitely at high zeta potentials. This behavior arises due to electrostatic charge condensation, finite ion size constraints, and nonlinear hydrodynamic interactions, all of which reduce the effective charge available for particle motion. As a result, classical theories systematically overestimate electrophoretic mobility in high-field regimes (Khair, 2022). Our study confirms that incorporating finite Debye length corrections, ion crowding effects, and viscoelastic fluid interactions is essential to accurately model electrophoresis in real-world applications.

Additionally, we demonstrated that nonlinear electrophoretic interactions influence colloidal stability and aggregation. In high-field environments, reduced electrophoretic repulsion enhances particle clustering, altering colloidal dispersion dynamics in biological, environmental, and microfluidic systems (Lozada-Cassou et al., 1999). These findings hold significant implications for contaminant transport, nanoparticle separation, and the design of electrophoretic devices.

By leveraging a multi-scale computational approach, our model successfully captures the transition from linear to nonlinear electrophoretic regimes. Numerical simulations using Finite Element Methods (FEM), spectral

techniques, and Monte Carlo simulations validate that nonlinear electrophoretic models provide more accurate predictions than traditional approximations. This work provides a comprehensive theoretical and computational framework for understanding electrophoretic motion beyond classical assumptions.

Open Problems & Future Directions

Despite the advancements made in this study, several challenges remain open for future investigation. One of the primary limitations of current electrophoretic models is their restriction to spherical colloidal particles. In reality, many biologically relevant molecules, industrial nanoparticles, and engineered microfluidic particles exhibit aspherical geometries, such as rods, ellipsoids, and platelets. The electrophoretic behavior of non-spherical particles is significantly more complex, as the interplay between hydrodynamic drag, electrostatic interactions, and particle orientation leads to anisotropic motion. Future studies must extend existing models to account for shape-dependent mobility variations, particularly in anisotropic colloidal suspensions (Felix et al., 2014).

Another critical challenge involves coupling electrophoresis with turbulent flow models in complex environments. Most electrophoretic studies assume laminar fluid motion, which is valid for small-scale systems and dilute suspensions. However, in realistic environmental conditions, such as rivers, oceans, and industrial reactors, fluid flow often exhibits turbulent behavior that interacts nonlinearly with charged particles. Understanding how turbulence modifies electrophoretic mobility, particle clustering, and contaminant transport remains an open question requiring advanced computational fluid dynamics (CFD) simulations and experimental validation (Hilton & Hayes, 2019).

Additionally, the role of multicomponent ionic solutions in electrophoresis is another frontier that warrants deeper exploration. Many real-world applications involve complex electrolytes containing multiple ionic species with varying valencies, hydration properties, and steric effects. The current study focuses on binary electrolytes, but future work should incorporate multi-ion electrokinetics, particularly for applications in biological electrophoresis, nanofluidics, and electrochemical separations (Barve et al., 2018). From a technological perspective, our findings suggest exciting opportunities for developing next-generation electrophoretic separation devices. The realization that mobility saturates in nonlinear regimes indicates that traditional electrophoretic separation methods might be inefficient at high field strengths. Future studies should explore how nonlinear mobility corrections can be leveraged for tunable microfluidic sorting, enabling precise control of particle separation based on charge, size, and shape (Tottori et al., 2019).

Furthermore, experimental validation remains a crucial aspect of advancing nonlinear electrophoretic models. While particle tracking velocimetry (PTV) and electrophoretic light scattering have provided valuable experimental benchmarks, high-resolution techniques such as optical tweezers, atomic force microscopy (AFM), and single-molecule electrophoresis should be employed to probe individual particle dynamics under extreme field conditions. These high-

precision experimental techniques can provide deeper insights into ion condensation, charge inversion, and electrohydrodynamic coupling, which remain poorly understood in strongly nonlinear electrophoretic systems (Lozada-Cassou et al., 1999).

In summary, this research establishes a solid theoretical and computational foundation for understanding nonlinear electrophoresis while highlighting several key challenges for future exploration. The extension to non-spherical particles, incorporation of turbulent flow interactions, multi-ion electrokinetics, and experimental validation are all crucial next steps toward developing a complete and predictive model for real-world electrophoretic phenomena. Addressing these challenges will not only refine our fundamental understanding of electrokinetics but also drive innovations in nanotechnology, environmental science, and biomedical diagnostics.

Final Remarks

This study represents a significant step forward in electrophoretic modeling, moving beyond classical approximations to capture essential nonlinear effects that dictate colloidal mobility, stability, and aggregation. By combining advanced numerical simulations, rigorous theoretical analysis, and experimental benchmarking, our findings bridge the gap between traditional electrophoretic theory and real-world applications. The insights gained from this research hold substantial potential for enhancing microfluidic technologies, optimizing environmental remediation strategies, and improving biomedical separation techniques.

As researchers continue to push the boundaries of nonlinear electrokinetics, addressing the open challenges outlined in this work will be instrumental in unlocking the full potential of electrophoretic transport in complex systems. Future investigations that integrate shape-dependent effects, turbulence interactions, multi-ion electrokinetics, and cutting-edge experimental validation will drive the next generation of advancements in this fundamental yet technologically critical field.

REFERENCES

- Barve, S., Babrekar, H., Bhave, T., & Bhoraskar, S. V. (2018). Segregation of nanoparticles by electrophoretic deposition technique: A mathematical model and its validation. *Powder Technology*, 327, 256-263. <https://www.sciencedirect.com/science/article/pii/S0032591018300457>
- Bhattacharyya, S., De, S., & Gopmandal, P. P. (2014). Electrophoresis of a colloidal particle embedded in electrolyte-saturated porous media. *Chemical Engineering Science*, 118, 188-198. <https://www.sciencedirect.com/science/article/pii/S0009250914003893>
- Felix, C., Yaroshchuk, A., Pasupathi, S., & Pollet, B. G. (2014). Electrophoresis and stability of nano-colloids: History, theory and experimental examples. *Advances in Colloid and Interface Science*, 211, 77-91. <https://www.sciencedirect.com/science/article/pii/S0001868614002000>
- Hilton, S. H., & Hayes, M. A. (2019). A mathematical model of dielectrophoretic data to connect measurements with cell properties. *Analytical and Bioanalytical Chemistry*, 411(12), 2565-2575. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6459731/pdf/nihms-1525067.pdf>
- Khair, A. S. (2022). Nonlinear electrophoresis of colloidal particles. *Current Opinion in Colloid & Interface Science*, 59, 101587. <https://www.sciencedirect.com/science/article/pii/S1359029422000267>
- Lobaskin, V., Dünweg, B., & Holm, C. (2004). Electrophoretic mobility of a charged colloidal particle: A computer simulation study. *Journal of Physics: Condensed Matter*, 16(38), S4063. <https://arxiv.org/pdf/cond-mat/0406278>
- Lozada-Cassou, M., González-Tovar, E., & Olivares, W. (1999). Nonlinear effects in the electrophoresis of a spherical colloidal particle. *Physical Review E*, 60(1), R17-R20. <https://pubmed.ncbi.nlm.nih.gov/11969842/>
- O'Brien, C. S. (2003). A mathematical model for colloidal aggregation. Thesis, University of Sydney. <https://core.ac.uk/download/pdf/154467026.pdf>
- Solomentsev, Y., Guelcher, S. A., Bevan, M., & Anderson, J. L. (2000). Aggregation dynamics for two particles during electrophoretic deposition under steady fields. *Langmuir*, 16(20), 9208-9216. <https://pubs.acs.org/doi/abs/10.1021/la0005199>
- Tottori, S., Misiunas, K., Keyser, U. F., & Bonthuis, D. J. (2019). Nonlinear electrophoresis of highly charged nonpolarizable particles. *Physical Review*

Letters, 123(1), 014502.
<https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.123.014502>